AUTHOR INDEX—VOLUME 20

Araya, J. E., J. E. Foster, and S. E. Cambron. A study of the biology of *Rhopalosiphum padi* (Homoptera: Aphididae) in winter wheat in northwestern Indiana. 47
Atkinson, Thomas H. (see Deyrup, Mark, and Thomas H. Atkinson) 59, 67
Barker, John F. Influence of temperature on egg production in a sunflower stem weevil *Cylindrocopturus adspersus* (Coleoptera: Curculionidae). 177
Bastian, R. A. (see Miller, F. D., Jr., et al.) 143
Calandrino, Frank J. (see Snider, Richard J., and Frank J. Calandrino) 1
Cambron, S. E. (see Araya, J. E., et al.) 47
Case, Laurie J. (see Jeffords, Michael R., and Laurie J. Case) 93
Cheetham, T. (see Miller, F. D., Jr., et al.) 143
Collins, Judith A., and Daniel T. Jennings. Spruce budworm and other lepidopterous prey of eumenid wasps (Hymenoptera: Eumenidae) in spruce-fir forests of Maine. 127
Cutkomp, L. K. (see Subramanyam, Bh., and L. K. Cutkomp) 99
Deyrup, Mark, and Thomas H. Atkinson. Comparative biology of temperate and subtropical bark and ambrosia beetles (Coleoptera: Scolytidae, Platypodidae) in Indiana and Florida. 59
Deyrup, Mark, and Thomas H. Atkinson. New distribution records of Scolytidae from Indiana and Florida. 67
Foster, J. E. (see Araya, J. E., et al.) 47
Fowler, Gary W., and Ann M. Lynch. Bibliography of sequential sampling plans in insect pest management based on Wald's Sequential Probability Ratio Test. 165
Fowler, Gary W., and Gary A. Simmons. Regression equations and table for estimating numbers of eggs in jack pine budworm (Lepidoptera: Tortricidae) egg masses in Michigan. 157
Fowler, Gary W., Bruce A. Montgomery, and Gary A. Simmons. Regression equations and table for estimating numbers of eggs in spruce budworm (Lepidoptera: Tortricidae) egg masses in Michigan. 161
Hart, E. R. (see Miller, F. D., Jr., et al.) 143
Heaton, George C. (see Wilson, Louis F., and George C. Heaton) 111
Hesch, P. G. (see Roeper, R. A., et al.) 173
Hilsenhoff, William L. An improved biotic index of organic stream pollution. 31
Hilsenhoff, William L. (see Lillie, R. A., et al.) 85
Hobrla, Sharon L. (see Wilson, Louis F., and Sharon L. Hobrla) 187
Jeffords, Michael R., and Laurie J. Case. Effect of prey density on diurnal activity and ovarian development in *Calosoma calidum* (Coleoptera: Carabidae): Implications for biological control of the gypsy moth, *Lymantria dispar* (Lepidoptera: Lymantriidae) in the Midwest. 93
Jennings, Daniel T., and Frank D. Parker. Habitats and spider prey of *Dipogon sayi sayi* (Hymenoptera: Pompilidae) in Washington County, Maine. 135
Jennings, Daniel T. (see Collins, Judith A., and Daniel T. Jennings) 127
Katovich, Steven A., and Herbert M. Kulman. *Leptoglossus corculus* and *Leptoglossus occidentalis* (Hemiptera: Coreidae) attacking red pine, *Pinus resinosa*, cones in Wisconsin and Minnesota. 119
Kirkendall, L. R. (see Roeper, R. A., et al.) 69
Korch, Peter P., III, and J. E. McPherson. Life history and laboratory rearing of *Gerris argenticollis* (Hemiptera: Gerridae) with descriptions of immature stages. 193
Kulman, Herbert M. (see Katovich, Steven A., and Herbert M. Kulman) 119
Kurczewski, Edmund J. (see Kurczewski, Frank E., and Edmund J. Kurczewski) 75, 81
Kurczewski, Frank E., and Edmund J. Kurczewski. Nest and prey of *Ageniella* (*Leucophrus*) *fulgifrons* (Hymenoptera: Pompilidae). 75
Kurczewski, Frank E., and Edmund J. Kurczewski. Northern distribution records for some Nearctic Pompilidae (Hymenoptera). 81
Kurczewski, Frank E., and Margery G. Spofford. Further observations on the nesting behavior of *Liris argentatus* (Hymenoptera: Sphecidae). 121

Larsen, C. D. (see Roeper, R. A., et al.) 173
Lillie, R. A., K. L. Schmude, and W. L. Hilsenhoff. Rediscovery of *Acanthametropus pecatonica* in the western Great Lakes region (Ephemeroptera: Siphlonuridae). 85
Lynch, Ann M. (see Fowler, Gary W., and Ann M. Lynch) 165
McPherson, J. E. (see Korch, Peter P., III and J. E. McPherson) 193
Miller, F. D., Jr., T. Cheetham, R. A. Bastian, and E. R. Hart. Parasites recovered from overwintering mimosa webworm, *Homadaula anisocentra* (Lepidoptera: Plutellidae). 143
Miller, William E. Change in nutritional quality of detached aspen and willow foliage used as insect food in the laboratory. 41
Montgomery, Bruce A. (see Fowler, Gary W., et al.) 161
Nielsen, David G. (see Purrington, Foster Forbes, and David G. Nielsen) 141
O'Brien, Mark F. Biology and distribution of *Tachysphex aethiops* in Michigan (Hymenoptera: Sphecidae: Larrinae). 71
Palik, B. J. (see Roeper, R. A., et al.) 69, 173
Parker, Frank D. (see Jennings, Daniel T., and Frank D. Parker) 135
Purrington, Foster Forbes, and David G. Nielsen. New host records of parasitic Hymenoptera in clearwing moths (Lepidoptera: Sesiidae). 141
Ragsdale, David W. (see Setiawan, Dwi P., and David W. Ragsdale) 103
Riotte, J. C. E. Designation of a neotype for *Sphinx poecila* Stephens (Lepidoptera: Sphingidae). 87
Rockett, C. Lee. Bacteria as ovipositional attractants for *Culex pipiens* (Diptera: Culicidae). 151
Roeper, R. A., D. V. Zestos, B. J. Palik, and L. R. Kirkendall. Distribution and host plants of *Corthylus punctatissimus* (Coleoptera: Scolytidae) in the Lower Peninsula of Michigan. 69
Roeper, R. A., B. J. Palik, D. V. Zestos, P. G. Hesch, and C. D. Larsen. Observations of the habits of *Corthylus punctatissimus* (Coleoptera: Scolytidae) infesting maple saplings in central Michigan. 173
Schmude, K. L. (see Lillie, R. A., et al.) 85
Schuder, Donald L. (see Yonker, James W., and Donald L. Schuder) 25.
Scriber, J. Mark. Puddling by female Florida tiger swallowtail butterflies, *Papilio glaucus australis* (Lepidoptera: Papilionidae). 21
Setiawan, Dwi P., and David W. Ragsdale. Use of aluminum-foil and oat-straw mulches for controlling aster leafhopper, *Macrosteles fascifrons* (Homoptera: Cicadellidae), and aster yellows in carrots. 103
Simmons, Gary A. (see Fowler, Gary W., and Gary A. Simmons) 157
Simmons, Gary A. (see Fowler, Gary W., et al.) 161
Snider, Richard J., and Frank J. Calandrino. An annotated list and new species descriptions of Collembola found in the Project ELF study area of Michigan. 1
Spofford, Margery G. (see Kurczewski, Frank E., and Margery G. Spofford) 121
Subramanyam, Bh., and L. K. Cutkomp. Total lipid and fatty acid composition in male and female larvae of Indian-meal moth and almond moth (Lepidoptera: Pyralidae). 99
Weitzner, Peter, and Mark E. Whalon. Head capsule widths as an indicator of the larval instar of codling moth (Lepidoptera: Olethreutidae). 147
Wellso, Stanley G., and Theo Wetzel. Economic thresholds for central European and North American wheat insects. 51
Wetzel, Theo (see Wellso, Stanley G., and Theo Wetzel) 51
Whalon, Mark E. (see Weitzner, Peter, and Mark E. Whalon) 147
Williams, Charles E. Exploitation of eggs of the Colorado potato beetle, *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae), by the exotic egg parasitoid *Edovum puttleri* (Hymenoptera: Eulophidae) in eggplant. 181
Wilson, Louis F., and George C. Heaton. Life history, damage, and gall development of the gall midge, *Neolasioptera brevis* (Diptera: Cicidomyiidae), injurious to honeylocust in Michigan. 111
Wilson, Louis F., and Sharon L. Hobrla. A procedure for sampling nymphs of Saratoga spittlebug, *Aphrophora saratogensis* (Homoptera: Cercopidae) using percentage of sample-units infested. 187
Yonker, James W., and Donald L. Schuder. Appearance of damage symptoms and reinfestation rates for Christmas trees attacked by the Zimmerman pine moth, *Dioryctria zimmermani* (Lepidoptera: Pyralidae). 25
Zestos, D. V. (see Roeper, R. A., et al.) 69, 173

INDEX TO ARTHROPOD TAXA—VOLUME 20

Ablabesmyia, 39
Acanthametropus pecatonica, 85–86
Acleris, 130–131
Acricotopus, 39
Acroneuria, 32, 37
Adelgid, pine leaf, 166
Aedes
  aegypti, 151
  atropalpus, 151
  communis, 151
  triseriatus, 151
Aeneolamia varia saccharina, 167
Aeshna, 37
Agapetus, 38
Agarodes distinctum, 38
Agathis cincta, 144
Agelenidae, 79
Ageniella
  accepta, 79
  agenioides, 78–79
  arcuata, 77–79
  argenteosignata, 75, 77–79
  blaisdelli, 75, 77–79
  conflicta, 75, 77–79
  coronata, 79
  cupida, 77
  euphorbiae, 79
  evansi, 79
  faceta, 79
  fulgifrons, 75–80, 82
  fuscipennis, 79
  longula, 78–79
  nivalis, 78–79
  norata, 77
  partita, 77–79, 82
  salti, 78–79
  semitincta, 77–79
  vogeli, 79
Agnetina capitata, 37
Agraylea, 38
Agrypnia, 33
Agriotes, 53, 55
Agromyza frontella, 167
Allocapnia, 32, 37
Allonemobius fasciatus, 123
Alloperla, 37
Ambrosiodmus
  devexulus, 65
  lecontei, 65
Ameletus, 37
Ammophila
  harti, 124
  urnaria, 75
Amphiagrion saucium, 37
Amphinemura
  delosa, 37
  linda, 37
Amphipoda, 39
Anabolia, 38
Anax junius, 37
Anceryx poecila, 87
Ancistrocerus, 139
  adiabatus, 127–133
  antilope, 127–133
Ancyronyx variegata, 38
Anoplius, 78
  cleora, 83
  depressipes, 83
  marginatus, 122
  nigerrimus, 83
  relativus, 83
Antestia, 167
Antestiopsis, 167
Antherea pernyi, 99
Antocha, 38
Anurida
  furcifera, 2
  granaria, 2
  pygmaea, 2
Anurophorus
  altus, 4
  binoculatus, 4
  septentrionalis, 4
Apanteles hyphantriae, 144
Aphid
  bird cherry-oat, 47–50
  cereal, 167
  green peach, 167
Aphididae, 47–50, 167
Aphrophora saratogensis, 187–192
Apis mellifera, 197
Aporinellus

taeniatus, 83
taeniolatus, 83
wheeleri, 83
Aporini, 82
Aporus
hirsutus, 82
niger, 82
Araneae, 135–139
Araptus sp., 68
Archips rosana, 130–131
Argia
apicalis, 37
moesta, 37
tibialis, 37
Argyrotaenia, 130–131
velutinana, 147
Armyworm, 53, 55
Arrhopalites
amarus, 10
benitus, 10
caecus, 10
Arthroplea bipunctata, 37
Arthropleona, 2–19
Asellus, 32
intermedius, 39
racovitzai, 39
Astata occidentalis, 77
Asynarchus montanus, 38
Atherix variegata, 38
Attaneuria ruralis, 37
Attenella attenuata, 37
Auplopodini, 82
Auplopus, 78
architectus, 82

Baetis
brunneicolor, 37
flavistriga, 37
frondalis, 37
hageni, 37
intercalaris, 37
longipalpus, 37
macdunnoughi, 37
propinquus, 37
pygmaeus, 37
vagans, 37
Baetisca
lacustris, 37
laurentina, 37
obesa, 37
Basiaeschna janata, 37
Beetle
cereal leaf, 52–53
Colorado potato, 181–186
Japanese, 167
June, 53, 55
Mexican bean, 167
mountain pine, 166
spruce, 166
Bembix, 71, 73
Bezzia, 38
Blepharicera tenuipes, 38
Blissus leucopterus 53, 55, 167
Bollworm, cotton, 167
Bombyx mori, 99
Borer
asiatic rice, 167
roundheaded, 166
Bothrosternini, 64
Bourletiella
dreisbachi, 11
hortensis, 10
koontzi, 10–17
russata, 10
Boyeria vinosa, 37
Brachycentrus
americanus, 38
incanus, 38
lateralis, 38
numerosus, 38
occidentalis, 38
Brachycerus, 37
Braconidae, 141, 144
Brillia, 33, 39
Budworm
jack pine, 157–160
spruce, 127–135, 161–163, 166
Bug
hairy chinch, 167
lygus, 167
Butterfly
cabbage, 99, 101
Florida tiger swallowtail, 21–23

Cabbageworm, imported, 167
Cadra cautella, 99–102
Caenis, 33, 37
Calicurgus hyalinatus, 82
Callibaetis, 37
Calopteryx
aequabilis, 37
maculata, 37
Calosoma
calidum, 93–97
frigidum, 94
scrutator, 94
sycophanta, 93–97
Camponotus pennsylvanicus, 197
Carabidae, 93–97, 183
Cardiocladius, 39
Carphoborus bifurcus, 64
Caterpillar

forest tent, 166
velvetbean, 167
Cecidomyiidae, 111–118
Centroptilum, 37
Cephus
cinctus, 53, 55
pygmaeus, 53, 55
Ceraclea, 38
Ceratopsyche, 31
alhedra, 38
alternans, 38
bronta, 38
morosa bifida, 38
morosa morosa, 38
recurvata, 38
riola, 38
slossonae, 38
sparna, 38
vexa, 38
walkeri, 38
Cerceris clypeata, 75
Cercopidae, 187–192
Ceropalinae, 83
Chaetocladius, 39
Chaoborus, 38
Chauliodes
pectinicornis, 38
rastricornis, 38
Cheumatopsyche, 33, 38
Chimarra
aterrima, 38
feria, 38
obscura, 38
socia, 38
Chirodamus fortis, 82
Chironomidae, 33, 39
Chironomus, 39
Choristoneura
fumiferana, 127–135, 161–163
pinus pinus, 157–160
rosaceana, 147
Choroterpes basalis, 37
Chramesus
chapuisi, 64
hicoriae, 64, 68
Chromagrion conditum, 37
Chrysogaster, 38
Chrysomelidae, 181–186
Chrysomelobia labidomerae, 181
Chrysopa, 38
carnea, 183
Chrysopidae, 38, 183
Cicadellidae, 103–109
Cladopelma, 39
Cladotanytarsus, 39
Clinotanypus, 39
Clioperla clio, 37
Cloeon alamance, 37
Cloverworm, green, 167
Clubionidae, 78–79
Cnephia dacotensis, 38
Cnesinus strigicollis, 64
Coccinellidae, 183
Coccotrypes distinctus, 65
Coenagrion resolutum, 37
Coleomegilla maculata, 183
Coleoptera, 33, 38, 52–53, 59–70, 93–97, 173–186
Coleotechnites, 130–131
Collembola, 1–19
Conchapelopia, 39
Conophthorus coniperda, 66
Contarinia
negundifolia, 118
tritici, 53–55
Cordites, 39
Cordulegaster maculatus, 37
Coreidae, 119–120
Corthylina, 66
Corthylini, 66
Corthylus
punctatisimus, 66, 68, 69–70, 173–175
spinifer, 66
Corydalus cornutus, 38
Corynoneura gracilis, 39
Cragonyx, 32
gracilis, 39
pseudogracilis, 39
Cricotopus, 39
Cryphalini, 61, 66
Cryptocarenus, 61
seriatus, 66
Cryptochironomus, 39
Cryptotendipes, 39
Cryptopygus decemoculatus, 4
Crypturgini, 65
Crypturgus
alutaceus, 65, 67
borealis, 67
pusillus, 67
Culex
pipiens pipiens, 151–155
quinquefasciatus, 151
tarsalis, 151, 167
Culicidae, 151–155
Culicoides, 38, 197
Culiseta inornata, 151
Curculionidae, 177–179
Cutworm, redbacked, 167
Cydia pomonella, 147–150
Cylindrocopturus adspersus, 177–179
Cyrnellus fraternus, 38

Dannella
lita, 37
simplex, 37
Delia coarctata, 53, 55
Demicryptochironomus, 39
Dendroctenus
terebrans, 64, 67
valens, 67
Diadegma, 144
Diamesa, 39
Dibrachus cavus, 141
Dicranoptycha winnemana, 197
Dicranota, 38
Dicrotendipes, 39
Dicyrtoma
atra, 12
aurata, 12
marmorata, 12
Dicyrtominae, 12
Didymops transversa, 37
Dioryctria zimmermani, 25–29
Diplectrona modesta, 38
Diplocladius, 39
Dipogon
calipterus, 137
sayi sayi, 135–139
Diptera, 33, 38–39, 53, 55, 121–125, 144, 151–155
Djalmabatista, 39
Dolichopodidae, 38
Dolophilodes distinctus, 38
Drosophila melanogaster, 195
Drunella cornutella, 37
Dryocoetes granicollis, 65
Dryocoetini, 65
Dubiraphia, 33
bivittata, 38
minima, 38
quadrinotata, 38
vittata, 38

Earworm, corn, 167
Ectemnia, 38
taeniatifrons, 197
Ectopria, 38
Edovum puttleri, 181–186
Einfeldia, 39
Elasmus albizziae, 143–145
Eleodes, 53, 55
Empididae, 38, 197
Empoasca assuta, 7
Enallagma
antennatum, 37
civile, 37
ebrium, 37
hageni, 37
Endochironomus, 33, 39
Entomobrya
assuta, 7
clitellaria, 7
comparata, 7
nivalis, 7
purpurascens, 7
Entomobryidae, 7–9
Entomobryinae, 7–8
Entypus unifasciatus, 75
Enytus obliteratus, 144
Epeorus vitreus, 37
Ephemera simulans, 37
Ephemerella
aurivillii, 37
catawba, 37
dorothea, 37
excrucians, 37
inermis, 37
invaria, 37
needhami, 37
species A, 33
subvaria, 37
Ephemeroptera, 32, 37, 85–86
Ephoron, 37
Ephydridae, 38
Episyron, 78
snowi, 82
Epitheca, 37
Epoicocladius, 39
Erioptera, 38
Eris sp., 76
Eristalis, 38
Eukiefferiella, 33, 39
Eulophidae, 118, 143–145, 181–186
Eumenidae, 127–133, 139
Euodynerus, 139
leucomelas, 127–133
Eurygaster integriceps, 167
Eurylophella
aestiva, 37
bicolor, 37
funeralis, 37
lutulenta, 37
temporalis, 37
Eurytoma, 118
Eurytomidae, 118
Eusimulium, 38
Evagetes
crassicornis, 82
subangulatus, 82
Evagora milleri, 166

Fleahopper, cotton, 167
Fly
Hessian, 53–54

wheat bulb, 53, 55
Folsomia
bisetosa, 4
nivalis, 4
Frenesia missa
Friesia sublimis, 2
Fruitworm, tomato, 167

Gammarus pseudolimnaeus, 39
Geocoris, 167
Gerridae, 193–204
Gerris
argentatus, 195
argenticollis, 193–204
buenoi, 193
marginatus, 195
najas, 197
pingreensis, 198
remigis, 198
Glossosoma, 38
Glyphopsyche irrorata, 38
Glyptotendipes, 39
Gnaphosidae, 79
Goera stylata, 38
Gomphurus, 37
Gomphus, 37
Grasshopper, 53, 55
Grub, white, 166–167
Gryllus
pennsylvanicus, 121–125
rubens, 123
veletis, 123–124
Guttipelopia, 39
Gymnopterus sp., 197

Habrophlebiodes americana, 37
Hagenius brevistylus, 37
Haplodiplosis equestris, 53
Haploperla brevis, 37
Harnischia, 39
Hastaperla, 37
Helichus
lithophilus, 38
striatus, 38
Helicopsyche borealis, 38
Heliothis, 167
Helius, 38
Hemiptera, 53, 55, 119–120, 193–204
Hemisturmia tortricis, 144
Heptagenia
diabasia, 37
flavescens, 37
hebe, 37
lucidipennis, 37
pulla, 37
Hesperoconopa, 38
Hesperophylax designatus, 38
Hetaerina americana, 37
Heterocloeon curiosum, 37
Heterotrissocladius, 39
Hexagenia, 37
Hexatoma, 38
Homadaula anisocentra, 143–145
Homoptera, 47–50, 53–54, 103–109, 187–192
Hyallela, 32
azteca, 39
Hydatophylax argus, 38
Hydrobaenus, 39
Hydropsyche
arinale, 38
betteni, 38
bidens, 38
cuanis, 38
dicantha, 38
leonardi, 38
orris, 38
phalerata, 38
placoda, 38
scalaris, 38
simulans, 38
Hydroptila, 38
Hylastes
exilis, 64
porculus, 67
salebrosus, 64
tenuis, 64, 67
Hylastini, 64
Hylastinus obscurus, 64
Hylema brassicae, 167
Hylesininae, 64
Hylesinini, 64
Hylesinus
aculeatus, 64
criddlei, 64
fasciatus, 64
pruinosus, 64
Hylocurus
bicornus, 67
binodatus, 65, 67
langstoni, 65
rudis, 65
torosus, 67
Hylogomphus brevis, 37
Hyloicus
plebia, 87
poecila, 87
Hylurgopinus rufipes, 64, 68
Hymenoptera, 53, 55, 71–84, 121–145, 181–186
Hypera postica, 167
Hypocryphalus mangiferae, 66
Hypogastrura nivicola, 2
Hypogastruridae, 2–3

Hypogastrurinae, 2
Hypothenemus, 60–61
  birmanus, 66
  brunneus, 66
  californicus, 66
  columbi, 66
  crudiae, 66
  dissimilis, 66
  eruditus, 66
  interstitialis, 66
  javanus, 66
  miles, 66
  rotundicollis, 66
  seriatus, 66
  sp., 66

Ichneumonidae, 141, 143–145
Ipini, 65
Ips
  avulsus, 65
  calligraphus, 65
  grandicollis, 65, 67
Ironoquia, 38
Ischnura verticalis, 37
Isogenoides
  frontalis, 37
  olivaceus, 37
Isonychia, 32, 37
Isoperla
  bilineata, 37
  clio, 37
  cotta, 37
  dicala, 37
  frisoni, 37
  lata, 37
  marlynia, 37
  nana, 37
  richardsoni, 37
  signata, 37
  slossonae, 37
  transmarina, 37
Isopoda, 39
Isotoma
  creli, 4
  japonica, 5
  nigrifrons, 4
  notabilis, 4
  nympha, 5–6
  viridis, 7
Isotomidae, 4–7
Isotomiella minor, 4

Katianninae, 10
Kiefferulus, 39

Labrundinia, 39
Larra, 123
Larrinae, 71–74
Larsia, 39
Leafhopper
  aster, 103–109
  potato, 167
Leafminer, alfalfa blotch, 167
Lebia grandis, 183
Lepidocyrtus
  helenae, 7
  hirtus, 8
  lignorum, 8
  paradoxus, 8
  violaceous, 8
Lepidoptera, 21–29, 38, 53, 55, 87–102, 127–133, 141–150, 157–163
Lepidostoma, 38
Leptinotarsa
  decemlineata, 181–186
  undecimlineata, 181
Leptoglossus
  corculus, 119–120
  occidentalis, 119–120
Leptophlebia, 37
Leucotrichia pictipes, 38
Leucrocuta hebe, 37
Leuctra
  ferruginea, 37
  sibleyi, 37
  tenella, 37
  tenuis, 37
Limnephilus, 38
Limnophila, 38
Limnophyes, 39
Limnoporus rufoscutellus, 198
Limonia, 38
Liparthrum squamosum, 67
Liris argentatus, 72, 121–125
Lissonota, 141
Litobrancha recurvata, 37
Looper
  cabbage, 167
  soybean, 94
Lycosidae, 78–79
Lymantria dispar, 93–97
Lymantor decipiens, 65
Lymantriidae, 93–97
Lype diversa, 38

Macrocentrus marginator, 141
Macronema, 38
Macronychus glabratus, 38
Macrosteles fascifrons, 103–109
Macrostemum zebratum, 38
Maggot, wheat stem, 53, 55
Malacasoma disstria, 166

Mayetiola
destructor, 53
rigidae, 118
Megaloptera, 38
Melanotus, 53, 55
Meromyza americana, 53, 55
Metopolophium dirhodum, 47, 53
Micracini, 65
Micracis
suturalis, 65
swainei, 65
Micracisella nanula, 65
Micrasema
kluane, 38
rusticum, 38
wataga, 38
Microcricotopus, 39
Microcylloepus pusillus, 38
Micropsectra, 39
Microtendipes, 39
Midge
lemon wheat blossom, 53–55
orange wheat blossom, 53–55
Miltogramminae, 121–125
Minagenia
congrua, 83
montisdorsa, 83
Minageniini, 83
Miogryllus verticallis, 123
Misumena vatia, 135–139
Molanna, 38
Monarthrum
fasciatum, 66
mali, 66
Monochamus, 166
Monodiamesa, 39
Monohelea, 38
Mosquito, northern house, 151–155
Moth
almond, 99–102
codling, 147–150
Douglas fir tussock, 166
gypsy, 93–97
Indian-meal, 99–102
Nantucket pine tip, 166
saddled prominent, 166
winter, 166
Zimmerman pine, 25–29
Muscidae, 38
Mystacides sepulchralis, 38

Nabis, 167
Nanocladius, 39
Natarsia, 39
Neanura muscorum, 3
Neanurinae, 2–3
Nectopsyche, 38
Needleminer, lodgepole, 165–166
Neelidae, 9
Neelus
incertus, 9
minimus, 9
minutus, 9
sniderí, 9
Nemotaulius hostilis, 38
Nemoura trispinosa, 37
Neodiprion swainei, 166
Neolasioptera brevis, 111–118
Neoperla stewarti, 37
Neophylax, 38
Neosminthurus bakeri, 11
Neotrichia, 38
Nephopteryx, 127–133
Neureclipsis, 38
Neurocordulia
molesta, 37
obsoleta, 37
yamakanensis, 37
Neuroptera, 183
Nigronia serricornis, 38
Nilobezzia, 38
Nilotanypus, 39
Nilothauma, 39
Nixe lucidipennis, 37
Nyctiophylax, 38
Nymphula, 38

Ochrotrichia, 38
Odonata, 33, 37
Odontella substriata, 2
Odontomesa, 39
Oecetis, 38
Oemopteryx glacialis, 37
Olethreutidae, 147–150
Oligostomis ocelligera, 38
Oligotrophus papyriferae, 118
Onocosmoecus quadrinotatus, 38
Onychiuridae, 3–4
Onychiurus
affinis, 3
allanae, 3
encarpatus, 3
parvicornis, 3
similis, 3
talus, 3
Operophtera brumata, 166
Ophiogomphus, 33, 37
Ophionellus foutsi, 144
Optioservus, 33, 38
fastiditus, 38
trivittatus, 38
Orchesella

ainsliei, 7
hexfasciata, 7
Orthocladius, 39
Orthoptera, 53, 55
Orthotomicus caelatus, 65
Orocharis saltator, 123
Oulema
lichinis, 53
melanopus, 52–53
Oxybelus bipunctatus, 124
Oxyethira, 38
Oxyopidae, 79

Pagastia, 39
Pagiocerus frontalis, 64
Palpomyia, 38
Papilio
glaucus australis, 21–22
glaucus canadensis, 21–22
glaucus glaucus, 21–22
polyxenes, 22
troilus, 22
Papilionidae, 21–23
Paracapnia angulata, 37
Parachironomus, 39
Paracladopelma, 39
Paragnetina media, 37
Paragyractis, 38
Paralauterborniella, 39
Paraleptophlebia, 37
Parametriochemus, 39
Parania geniculata, 143–145
Paraolinx canadensis, 144
Paraphaenocladius, 39
Parapoynx, 38
Parapsyche apicalis, 38
Paratanytarsus, 39
Paratendipes, 39
Pedicia, 38
Pediobius, 118
Pentagenia vittigera, 37, 86
Pentaneura, 39
Pepsinae, 82
Pepsini, 82
Pericoma, 38
Perlesta placida, 32, 37
Perlinella
drymo, 37
ephyre, 37
Petrophila, 38
Phaenopsectra, 33, 39
Phaeogenes ater, 141
Phanagenia, 78
bombycina, 77
Phasganophora, 37
Phidippus audax, 75
Philanthus
gibbosus, 77
politus, 122
Phloeosinini, 64
Phloeosinus dentatus, 64, 68
Phloeotribini, 64
Phloeotribus
dentifrons, 64
frontalis, 64, 68
liminaris, 64
Phrosinella aurifacies, 122
Phyrganea, 38
Phylocentropus placidus, 38
Phyllophaga, 53, 55
Pieris
brassicae, 99, 101
rapae, 167
Pilaria, 38
Pinworm, tomato, 167
Pityoborus comatus, 66
Pityogenes
hopkinsi, 65
plagiatus, 68
Pityophthorina, 66
Pityophthorus
annecteus, 66
cariniceps, 68
confusus, 66
crinalis, 66
lautus, 66
liquidambarus, 68
opaculus, 66
pinivorus, 66
puberulus, 66
pulchellus, 68
pulicarius, 66
pullus, 68
sp., 66
Plathemis lydia, 37
Platycentropus, 38
Platygaster, 118
Platygasteridae, 118
Platypodidae, 59–64
Platypus
compositus, 64
flavicornis, 64
parallelus, 64
Plecoptera, 32, 37
Plodia interpunctella, 99–102
Plutellidae, 143–145
Podalonia luctuosa, 72, 124
Podosesis
aureocincta, 141
syringae, 141
Poecilopompilus interruptus, 83
Polycentropus, 38

Polygraphini, 64
Polypedilum, 39
Pompilidae, 75–84, 135–139
Pompilinae, 82–83
Pompilini, 82–83
Potamanthus, 37
Potamyia flava, 38
Potthastia, 39
Premnobius cavipennis, 65
Priocnemis
  aequalis, 82
  notha, 82
Probezzia, 38
Procladius, 39
Prodiamesa, 39
Proisotoma minima, 4
Prosimulium
  fuscum, 38
  gibsoni, 38
  magnum, 38
  mixtum, 38
  mysticum, 38
Prostoia completa, 37
Protoptila, 38
Psectrocladius, 39
Psectrotanypus, 39
Pseudachorutes
  aureofasciatus, 2
  indiana, 2
  saxatilis, 2
Pseudaletia unipunctata, 53, 55
Pseudatomoscelis seriatus, 167
Pseudiron centralis, 37, 86
Pseudochironomus, 39
Pseudocloeon
  carolina, 37
  dubium, 37
  myrsum, 37
  parvulum, 37
  punctiventris, 37
  species A, 33, 37
Pseudolimnophila, 38
Pseudomittia, 39
Pseudopityophthorus
  asperulus, 66
  minutissimus, 66
Pseudoplusia includens, 94
Pseudorthocladius, 39
Pseudosinella
  rolfsi, 8
  violenta, 8
Pseudostenophylax uniformis, 38
Pseudothysanoes
  lecontei, 65
  rigidus, 65
Pseuphenus herricki, 38
Psiloptreta indecisa, 38
Psychoda, 38
Psychoglypha subborealis, 38
Psychomyia flavida, 38
Pteromalidae, 141
Pteronarcys, 32, 37
Ptilostomis, 38
Pycnopsyche, 38
Pyralidae, 25–29, 99–102, 127–133

Recurvaria milleri, 165–166
Rheocricotopus, 39
Rheotanytarsus, 39
Rhithrogena, 37
Rhopalosiphum
  insertum, 53
  maidis, 53
  padi, 47–50, 53
Rhyacophila
  brunnea, 38
  fuscula, 38
  lobifera, 38
  vibox, 38
Rootworm, corn, 167

Saetheria, 39
Salticidae, 75–79
Sarcophagidae, 121–125
Sawfly
  European wheat stem, 53, 55
  jack pine, 166
  red pine, 166
  hemlock, 166
  larch, 166
  wheat stem, 53, 55
Schizaphis graminum, 53–54
Scolytidae, 59–70, 173–175
Scolytinae, 65
Scolytini, 65
Scolytus
  mali, 65
  multistriatus, 65
  muticus, 65
  quadrispinosus, 65, 68
Senotainia
  trilineata, 121–125
  vigilans, 124
Serratella
  deficiens, 37
  sordida, 37
Sesiidae, 141–142
Setodes, 38
Shipsa rotunda, 37
Sialis, 38
Silkworm, 99
  oak, 99

Simulium
aurium, 39
corbis, 39
croxtoni, 39
euryadminiculum, 39
fibrinflatum, 39
jenningsi, 39
johannseni, 39
latipes, 39
luggeri, 39
pictipes, 39
rugglesi, 39
tuberosum, 39
venustum, 39
verecundum, 39
vittatum, 31, 33, 39
Siphlonuridae, 85–86
Siphlonurus, 37
Siphloplecton, 37
Siphluriscus chinensis, 85
Sitobion
avenae, 47, 53–54
fragariae, 53
Sitodiplosis mosellana, 53–55
Sminthuridae, 9
Sminthurides
lepus, 9
occultus, 9
pumilis, 9
Sminthuridinae, 9–10
Sminthurinae, 10–18
Sminthurinus
henshawi, 10
intermedius, 10
macgillivrayi, 10
quadrimaculatus, 10
Sminthurus
butcheri, 12
nigromaculatus, 12, 18
purpurescens, 12, 18
Smittia, 39
Somatochlora, 37
Soyedina vallicularia, 37
Sphaeromais longipennis, 39
Spharagemon sp., 71–73
Sphecidae, 71–74, 121–125
Sphecius speciosus, 122–123
Sphingidae, 87–91
Sphinx
eremitus, 87
gordius, 87
lineolata, 87
pinastri, 87
poecila, 87–91
Spilonota ocellana, 130–131
Spinadis, 37, 86
Spittlebug, Saratoga, 187–192
Stactobiella, 38
Stegopterna mutata, 39
Stempellina, 39
Stempellinella, 39
Stenacron interpunctatum, 37
Stenelmis, 33, 38
crenata, 33
Stenochironomus, 39
Stenonema
exiguum, 37
femoratum, 37
integrum, 37
mediopunctatum, 37
mexicanum, 37
modestum, 37
pulchellum, 37
terminatum, 37
vicarium, 37
Stictochironomus, 39
Strophopteryx fasciata, 37
Stylogomphus albistylus, 37
Stylurus, 37
Sympetrum, 37
Symphitopsyche, 38
Symphypleona, 9–18
Sympotthastia, 39
Synanthedon pictipes, 141
Synorthocladius, 39

Tabanus, 39
Tachinidae, 144
Tachysphex
aethiops, 71–74
pompiliformis, 73
punctifrons, 73
tarsatus, 73
Tachytes
intermedius, 123
mergus, 123
Taeniopteryx, 32, 37
Tanypus, 39
Tanytarsus, 39
Thienemanniella, 39
Thomisidae, 79
Thysanoes fimbricornis, 65
Tipula, 39
Tomicini, 64
Tomocerinae, 9
Tomocerus
flavescens, 9
lamelliferus, 9
Tortricidae, 127–133, 157–163
Triaenodes, 38
Tribelos, 33, 39
Trichoptera, 33, 38

Tricorythodes, 37
Trimerotropis sp., 71, 73
Trischidias
  atoma, 66
  exigua, 66
  sp., 66
Tullbergia
  clavata, 3
  granulata, 3
  malla, 4
  yosiii, 4
Tvetenia, 33, 39

Webworm, mimosa, 143–145
Weevil
  alfalfa, 167
  boll, 167
Willemia
  denisi, 2
  intermedia, 2
  similis, 2
Willowsia buski, 7
Wireworm, 53, 55
  southern potato, 167
Wormaldia moestus, 38

Xenochironomus, 39
Xenylla
  acauda, 2
  pallescens, 2
Xyleborini, 65–66
Xyleborinus saxeseni, 65
Xyleborus, 60
  affinus, 65
  celsus, 65
  dispar, 65
  ferrugineus, 65
  obesus, 65
  planicollis, 65, 68
  pubescens, 65
  sayi, 65
  tachygraphus, 68
  volvulus, 65
  xylographus, 65
Xylosandrus
  compactus, 66
  crassiusculus, 66, 68
  germanus, 66
Xyloterinus politus, 65
Xyloternini, 65
Xylotopus par, 33, 39
Xysticus
  emertoni, 135–139
  ferox, 138
  punctatus, 135–139

Zabrus tenebroides, 53
Zalutschia, 39
Zavrelimyia, 39